

Marius-Nicușor Grigore
Constantin Toma

HALOFITELE

ASPECTE DE ANATOMIE ECOLOGICĂ



Editura Universității „Alexandru Ioan Cuza” Iași

MARIUS-NICUȘOR GRIGORE

CONSTANTIN TOMA

HALOFITELE
Aspecte de anatomie ecologică

Referenți științifici:

Dr. Monica Tereza Boșcaiu Neagu

Universidad Politecnica de Valencia

Prof. univ. dr. Maria-Magdalena Zamfirache

Universitatea „Alexandru Ioan Cuza” Iași, Facultatea de Biologie

Conf. univ. dr. Lăcrămioara Ivănescu

Universitatea „Alexandru Ioan Cuza” Iași, Facultatea de Biologie

Redactor: Oana Bilan

Tehnoredactare: Marius-Nicușor Grigore, Luminița Răducanu

Coperta: Manuela Oboroceanu

(pe copertă: *Camphorosma annua* – secțiuni transversale prin limbul foliar și
Aster tripolium – vedere de ansamblu)

ISBN 978-973-703-542-4

© Editura Universității „Alexandru Ioan Cuza”, 2010

700109 – Iași, str. Pinului, nr. 1A, tel./fax: (0232) 314947

[http:// www.editura.uaic.ro](http://www.editura.uaic.ro) e-mail: editura@uaic.ro

MARIUS-NICUȘOR GRIGORE

CONSTANTIN TOMA

HALOFITELE

Aspecte de anatomie ecologică

Cuvânt înainte de T. J. Flowers



Editura Universității „Alexandru Ioan Cuza” Iași
2010

Descrierea CIP a Bibliotecii Naționale a României
GRIGORE, MARIUS-NICUȘOR

Halofitele : aspecte de anatomie ecologică / Marius-Nicușor
Grigore, Constantin Toma ; cuv. înainte de Timothy Flowers. - Iași :
Editura Universității "Alexandru Ioan Cuza", 2010

Bibliogr.

ISBN 978-973-703-542-4

I. Toma, Constantin, biologie
II. Flowers, Timothy (pref.)

581.526.52

CUPRINS

FOREWORD	7
CUVÂNT ÎNAINTE	9
INTRODUCERE	11

PARTEA I

CONSIDERAȚII TEORETICE

1. Definiția și clasificarea halofitelor. Dificultăți inerente	13
2. Halofitele și mediile lor de viață. Locul halofitelor în clasificarea ecologică a plantelor	26
3. Adaptări morfo-anatomice generale ale halofitelor	30
4. Halofitele și stresul salin. Trăsăturile histo-anatomice ale halofitelor: efect formativ, adaptare sau răspuns la un factor abiotic exagerat?	34

PARTEA A II-A

STUDIUL HISTO-ANATOMIC AL UNOR SPECII DE HALOFITE DIN FLORA MOLDOVEI

1. Material și metodă de lucru	37
2. Caracterizarea ecologică și clasificarea halofitelor luate în studiu	40
3. Rezultate și discuții	43
<i>Artemisia santonicum</i> L.	43
<i>Aster tripolium</i> L. ssp. <i>pannonicus</i> (Jacq.) Soo	54
<i>Lactuca saligna</i> L.	69
<i>Scorzonera cana</i> (C. A. Mey.) O. Hoffm.	80
<i>Lepidium cartilagineum</i> (J. C. Mayer) Thell. ssp. <i>crassifolium</i> (Waldst. & Kit.) Thell. ...	92
<i>Lepidium latifolium</i> L.	100
<i>Lepidium perfoliatum</i> L.	109
<i>Spergularia media</i> (L.) C. Presl	117
<i>Atriplex littoralis</i> L.	126
<i>Atriplex prostrata</i> Boucher ex DC	139
<i>Atriplex tatarica</i> L.	152
<i>Bassia hirsuta</i> (L.) Asch.	163
<i>Camphorosma annua</i> Pall.	172
<i>Halimione verrucifera</i> (M. Bieb.) Aellen	180
<i>Petrosimonia oppositifolia</i> (Pall.) Litv.	193

<i>Petrosimonia triandra</i> (Pall.) Simonk.	202
<i>Salicornia europaea</i> L.	211
<i>Suaeda maritima</i> (L.) Dumort.	223
<i>Bolboschoenus maritimus</i> (L.) Palla ssp. <i>compactus</i> (Hoffm.) Drobow	229
<i>Carex distans</i> L.	234
<i>Iris halophila</i> Pall.	241
<i>Juncus gerardi</i> Loisel.	250
<i>Trifolium fragiferum</i> L.	261
<i>Puccinellia distans</i> (L.) Parl. ssp. <i>limosa</i> (Schur) Jáv.	269
<i>Plantago schwarzenbergiana</i> Schur	273
<i>Polygonum patulum</i> M. Bieb.	284
 CONCLUZII	 293
 ENGLISH ABSTRACT	 300
 BIBLIOGRAFIE	 304

ABSTRACT

Salinisation is currently one of the biggest problems facing Earth agriculture. In a continuous increase of world population, obtaining of salt tolerant crops represents a true priority. Halophytes have been defined as salt-tolerant plants having therefore, cellular, biochemical, physiological and anatomical mechanisms allowing them to survive in high salinity conditions.

This work represents an anatomical and ecological monograph of halophyte species from Moldova (North-Eastern Romania), representing the results of investigations conducted over many years. Thus, 26 taxa were investigated; these species belong to different botanical families and constitute a heterogeneous group of plants collected from different saline – some dry, others wet habitats.

Ecological anatomy approach involves a correlation between structure and habitat, between certain well-defined anatomical adaptations and some environmental factors such as salinity, soil moisture or dryness. In some cases, finding such correlations has been proved facile, while in other cases, some anatomical adaptations remain controversial, because their function is not fully elucidated.

In the introductory part of the book, we have discussed some theoretical issues related to definition and classification of halophytes and the relationship between halophytes and salt stress, the latter one being one of the most intensely studied in plant biology field. We are keeping open the question whether anatomical adaptations of halophytes are the result of formative effect of salts, salt stress damage or an expression of positive adaptation, allowing plants to withstand in hypersaline environments.

The proper part of the work refers to the histo-anatomical study of halophyte species. Ecological interpretations were made whenever possible, in concordance with the evidenced and discussed structure. As a general rule, we have seen that in most cases, histo-anatomical features are logically linked to environmental factors, thus being the effect of environmental factors.

We found that taxa included in the *Chenopodiaceae* (*Atriplex littoralis*, *A. prostrata*, *A. tatarica*, *Camphorosma annua*, *Halimione verrucifera*, *Petrosimonia oppositifolia*, *P. triandra*, *Salicornia europaea*, *Suaeda maritima*) show successive cambia phenomenon on vegetative axial organs (root, stem) level. Regarded by many authors as a structural anomaly, the connection of this phenomenon with salinity factor remains unclear. Although the interpretation of this phenomenon is very difficult to be done in ecological and adaptive direction, because here is almost a

completely lack in some studies, so we tried, as a hypothesis, to explain this phenomenon. We have assumed that additional cambia phenomenon might be a mechanism involved in regulation of salt content, thus having an adaptive and ecologic significance. Certainly, it has been sometimes shown the fact that the intense lignification which is provided by the activity of successive cambia can be stimulated by salinity. Therefore, this could confer mechanical resistance of cells to high osmotic pressure of soil solution.

Halophyte species collected from wet habitats (*Aster tripolium* ssp. *pannonicus*, *Juncus grardi*, *Spergularia media*) present aerenchyma, especially in underground organs (rhizome, root). This feature is a common adaptation in species vegetating in such habitats, supplying the plant with oxygen during the possible hypoxia conditions.

Another interesting anatomical feature is a well developed endodermis, evidenced in some species (*Artemisia santonicum*, *Juncus gerardi*, *Lactuca saligna*, *Salicornia europaea*, *Suaeda maritima*, *Bolboschoenus maritimus*), on axial vegetative organs level. We have viewed this structure as a barrier that controls and limits the salt penetration in plant body.

Other adaptation, relatively common on halophytes and intensely discussed, is the shoot succulence. We have evidenced this feature, expressed by a water storage tissue and sometimes by a well-developed palisade tissue, in: *Salicornia europaea*, *Suaeda maritima*, *Petrosimonia oppositifolia*, *P. triandra* and *Spergularia media*. On halophyte species growing in wet habitats (*Suaeda*, *Salicornia*), the succulence ensures the erect position of plant and the potential dilution of toxic salts. In addition, the succulence can be induced and developed by chlorine ions as it is known that the two species listed above have been included in the chloruric plant associations. In xero-halophytes (*Petrosimonia oppositifolia* and *P. triandra*) the water storage tissues constitute, in the same time, a water reservoir needed in the dry seasons. Moreover, at these species, the succulence also facilitates the salt dilution, because in dry conditions salts become more concentrated in root zone.

Traheids occurring in succulent tissues of *Salicornia* are very interesting but also controversial structures. It is assumed that these tracheids are involved in plant water balance, salt tolerance, mechanical support and water or air storage. They are located among the palisade cells, approximately perpendicular to the epidermis, with no direct connection neither with epidermis, on the one hand, neither with vascular system of plant, on the other hand.

The vezicular, salt-secreting hairs (bladders) of *Atriplex tatarica* and *Halimione verrucifera*, evidenced in the leaf structure, are highly specialized structures, involved in the accumulation and subsequent removal of salts in excess. They are characteristic mainly in the *Chenopodiaceae*, and consist in a stalk-cell and a vezicular (bladder) cell, where salts are accumulated in order to be eliminated on leaf surface. The secretory pathway, chosen by many halophyte species (crynohalophytes) represents an important adaptive strategy in salt tolerant plants life.

Another adaptive mechanism related to the C₄ photosynthesis, is the so-called Kranz Anatomy configuration, in the leaf structure of some *Chenopodiaceae* species. Thus, we have described the atriplicoid sub-type on *Atriplex tatarica*, meaning that here is a sheath formed by izodiametric cells that form a layer more or less continuous around the vascular bundles. On *Petrosimonia opositifolia*, *P. triandra* and *Camphorosma annua* we have described the kochioid sub-type included in Kranz anatomy. This means, mainly, the following tissues disposition: epidermis (with hypodermis on *Camphorosma*), external chlorenchyma (palisade tissue), internal clorenchyma (a layer of more or less cubic cells) and central, water storage tissue in which usually here a great vascular bundle is located. The two chlorenchyma tissues are arranged in two discontinuous layers, around the leaf.

We have also evidenced bulliform cells (*Puccinellia distans*, *Juncus gerardi*, *Bolboschoenus maritimus*, *Carex distans*), at leaf epidermis level; these cells are mostly involved in leaves rolling in dry habitat conditions. But, apparently, their role is more complex and subtle than at first sight. Although they are considered a xerophytic character, we have evidenced them on hygro-halophyte species. In such a context, we think that here is not relevant to consider one single environmental factor, seen stable and fixed, but its fluctuation during one growing season. The plants listed above have been called amphibious halophytes, due to their ability to survive in dry soil conditions, and this real possibility requires, in addition, the presence of xeromorphytic features.

In general, as a result of our investigations, we can say that there are some obvious anatomical adaptations of halophytes, well defined, enabling them to survive in hypersaline environments. These features characterize the „obligatory” halophytes – species adapted to high salinity conditions, which are strictly restricted to salty areas, and are not found on other soil types than the saline ones. These observations allowed us to formulate a hypothesis regarding the „extreme” adaptations of halophytes to salinity. The most evident and „spectacular” characters, linked to halophilous profile

were observed on halophyte species adapted to higher salinity conditions (and usually on the species that occur only in high salinized areas).

Following our investigations and given interpretations, we have proposed a new system of halophytes classification, based on the relevance of anatomical adaptations correlated with the intensity of environmental factors (most oftenly, soil salinity). Thus, the *Chenopodiaceae* succulent species best adapted to high salinity conditions (*Salicornia*, *Suaeda*, *Halimione*, *Petrosimonia oppositifolia* and *P. triandra*) were nominated as extremohalophytes; in addition, because they grow only in very salinized environments, we applied to them the term irreversible halophytes.

Atriplex littoralis, *A. prostrata*, *A. tatarica*, *Bassia hirsuta*, *Camphorosma annua* species, which have a large ecological spectrum and they are not so strictly related to increased salinity, were called by us as reversible halophytes. These species may also pass in less salinized environments.

The supporting, accidental or preferential halophytes which don't have histo-anatomical characters very clearly correlated to salinity factor were called mesohalophytes. They are more subjected to multiple influences of environmental factors, including soil salinity, but this is not the major factor inducing spectacular adaptations within this group of halophyte species. In the mesohalophytes group, the above mentioned amphibious halophytes could be included.

Our investigations and observations suggest that here is a relevant correlation between histo-anatomical features of halophytes and ecological factors of environments where they vegetate.

Marius - Nicușor Grigore
Constantin Toma

HALOFITELE

ASPECTE DE ANATOMIE ECOLOGICĂ



Halofitele alcătuiesc un grup ecologic de plante, ce prezintă interesante adaptări anatomice – unele dintre ele mult discutate și controversate în legătură cu factorul salinitate.

În acest context, prezenta monografie constituie o combinație inedită datorită modalității de abordare. Pe de o parte, este o lucrare de fitoanatomie, în sensul ideii fundamentate și individualizate de Școala de Anatomie Vegetală ieșeană, iar pe de altă parte, o lucrare care depășește cu mult simplele descrieri anatomice. Cartea are modesta, dar îndrăzneța intenție de a promova un gen de anatomie modernă, care să rămână tangențială cu preocupările interdisciplinare de ecologie, ecofiziologie și implicații ale stresului salin – o tendință generală care domină momentan climatul științific mondial al biologiei vegetale.

www.editura.uaic.ro



9 789737 035424

Editura Universității „Alexandru Ioan Cuza” Iași